



TAXONOMIC STUDY OF TWO GENUS IN LICE
***Linognathus Africanus* (Kellogg And Paine, 1911) &**
***Bovicola (Damalinia) ovis* (Schrank, 1781) BY SCANNING**
ELECTRONIC MICROSCOPE (SEM)

[3]

Sara A. Aljubran¹ and Souad M. Alsaqabi^{1 a)}

1- University of Dammam, College of Science, Department of Biology, Section Zoology

*(Mailing Address: P.O. Box 838 Postal Code 31113, University of Dammam, Dammam, Saudi Arabia.

a) E-Mail: Dr-Alsaqabi@hotmail.com

Keywords: *Bovicola (Damalinia) ovis*,
Linognathus africanus, Saudi Arabia,
SEM.

ABSTRACT

This study presents the Mallophaga and Anoplura lice which affects farm animals (camels sheep and goats), a local area east of the Saudi Arabia of different ages, races, classes. Study the classification of

lice were recorded as using the optical microscope and electronic scanning. The study shows scanning electron microscopy micro-structure of the species: *Bovicola (Damalinia) ovis*, *Linognathus africanus* and the study showed the installation of a portfolio of head lice, and the distribution of filament head, chest and abdomen, as shown over the centuries, the difference in the form of remote sensing and Swallows in thesexuallice, the first study in this area.

(Received September 29, 2010)
(Accepted January 17, 2011)

تحكيم: أ.د خالد عبد العزيز عبد العاطى